



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

never heard before, even in mild seasons, of *C. auratus* wintering so far north. — J. B. WILLIAMS, *Montreal, Can.*

Chuck-will's-widow in Kansas. — Prof. D. E. Lantz kindly informs me that the specimen of the Chuck-will's-widow (*Antrostomus carolinensis*) taken at Wichita, Kan., as recorded by me in a footnote to p. 187 of 'The Auk' for April, 1889, is really the specimen recorded by him on the same page, which was sent to Prof. Dyche for confirmation of the identification, and by the latter forwarded to me. — J. A. ALLEN, *Am. Mus. Nat. History, New York City.*

Otocoris alpestris praticola at Ipswich, Mass. — On October 26, 1899, at Ipswich, Mass., with Dr. Walter Faxon and Mr. G. M. Allen, I took a male Prairie Horned Lark out of a flock of four birds (two others were also seen later), the other three appearing to be of the same race. The specimen taken seems of especial interest, as its measurements and paleness make it approach very nearly to *arenicola*, — the specimen being almost intermediate as it is. Mr. Harry C. Oberholser and Mr. William Brewster, and all who have examined the bird, are of the opinion, however, that it is perhaps nearest *praticola*, and must be called such. Geographically of course, the bird is *praticola*, for if *arenicola* it is a straggler far out of its usual range. — REGINALD HEBER HOWE, JR., *Longwood, Mass.*

The Red Crossbill Unusually Common at Portland, Maine, in Summer. — During the early summer of 1899, Red Crossbills (*Loxia curvirostra minor*) were numerous about the west end of Portland, Maine, — a section of the city in which there is still a good deal of land devoted to gardens and lawns, and which contains a hill-side park with a large grove of pines. I returned to Portland from the south on June 2. The birds were then everywhere in evidence about my neighborhood, but mainly because of their vociferousness. They were quite elusive, and I found it difficult to estimate their numbers except from the noise they made. It was not until July 1 that I saw a large number together. On that morning, I counted twenty-eight in one open flock which flew slowly over me as I stood in the middle of a wide lawn. On the afternoon of that same day I left Portland for the summer. — NATHAN CLIFFORD BROWN, *Portland, Me.*

White-winged Crossbills in Rhode Island. — It may be of interest to Rhode Island ornithologists to record the taking of White-winged Crossbills (*Loxia leucoptera*) at Neutaconkanut Hill, Johnston, R. I., Jan. 14 and 30, and at Pawtuxet on the 1st, 2d and 24th of February. — EDWARD H. ARMSTRONG, *Providence, R. I.*

White-winged Crossbills and Brünnich's Murres in Central New Hamp-